


Smooth Manifolds and Their Tangent Bundles

Tangent bundles to smooth manifolds are one of the oldest and most important examples of vector bundles.

Def'n: A smooth structure on a d -dim'l mfld

M^d is a collection of charts $h_i: U_i \rightarrow M^d$

(U_i open in \mathbb{R}^d) whose images cover M^d and whose transition functions $h_j^{-1} \circ h_i$ are smooth.



- Note that $h_j^{-1} \circ h_i$ is defined on $h_i^{-1}(h_j(U_j) \cap h_i(U_i))$.

- We call a map $h: U \rightarrow V$ ($U \subset \mathbb{R}^n$, $V \subset \mathbb{R}^m$ open sets) smooth if all partial derivatives of all orders exist and are continuous.

Tangent bundles to smooth manifolds can be described abstractly (see Spivak, Diff. Geom. Vol. I), but it is much easier to define tangent bundles to submanifolds of \mathbb{R}^n .

because we already know how to define tangent vectors in \mathbb{R}^n :

Following Milnor-Stasheff, we have:

Defn: A subset $M^d \subseteq \mathbb{R}^N$ is a (sub)-manifold of dimension d if at each point $x \in M^d$ there exists an open set $U \in \mathbb{R}^d$ and a smooth map $h: U \rightarrow M^d$ (i.e. $U \xrightarrow{h} M^d \hookrightarrow \mathbb{R}^N$ is smooth)

such that

- 1) h is a homeomorphism onto its image;
- 2) the vectors $\frac{\partial h}{\partial u_j} = \begin{pmatrix} \frac{\partial h_1}{\partial u_j} \\ \vdots \\ \frac{\partial h_N}{\partial u_j} \end{pmatrix} \in \mathbb{R}^N$ ($j=1, \dots, d$) are linearly independent

(i.e. the Jacobian $\left(\frac{\partial h_i}{\partial u_j}(a)\right)$ has maximum rank).

[We call h a local parameterization.]

Remarks - Condition 1) implies that M^d is a topological mfld.

- Condition 2) implies that M^d is a smooth mfld:
Using the Implicit Function theorem one can check that the transition funcs g_{ijh} are automatically smooth.
(Lemma 1.1 in Milnor-Stasheff).

- The notation $\frac{\partial h_i}{\partial u_j}$ means we write $h(a) = (h_1(a), \dots, h_N(a))$ $a \in \mathbb{R}^d$
and then $\frac{\partial h_i(a)}{\partial u_j} = \lim_{t \rightarrow 0} \frac{h_i(a_1, \dots, a_{j-1}, t, a_{j+1}, \dots, a_N) - h_i(a)}{t}$
 $(\bar{a} = (a_1, \dots, a_d))$

We now want to define tangent vectors to points $x \in M$. These arise as velocity vectors to smooth paths through x .

Defn: If $\alpha: [a, b] \xrightarrow{\text{SR}} M^d \subseteq \mathbb{R}^N$ is smooth and $\alpha(t_0) = x$, then the velocity vector of α at t_0 is $\alpha'(t_0) = \left(\frac{d\alpha_1}{dt}(t_0), \dots, \frac{d\alpha_N}{dt}(t_0) \right)$.

Def'n: The tangent space $T_x M^d$ to $M^d \subseteq \mathbb{R}^N$ at a

point $x \in M^d$ is $\{\vec{v} \in \mathbb{R}^N \mid \vec{v} = \alpha'(t) \text{ for some smooth path } \alpha: \mathbb{R} \rightarrow M^d \text{ w/ } \alpha(t) = x\}$.


Lemma 1.2: Say $h: U \rightarrow M^d$ is a local parametrization of M^d , with $x = h(\bar{u})$. Then

$$T_x M^d = \text{Span}\left(\frac{\partial h}{\partial u_1}(\bar{u}), \dots, \frac{\partial h}{\partial u_d}(\bar{u})\right).$$

Proof: For any $\sum_{i=1}^d c_i \frac{\partial h}{\partial u_i}(\bar{u}) \in \text{Span}\left(\frac{\partial h}{\partial u_i}(\bar{u})\right)$, there

is a line $\alpha(t) = \bar{u} + t \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_d \end{pmatrix}$ ($\alpha: \mathbb{R} \rightarrow \mathbb{R}^d$) such that


$$(h \circ \alpha)'(0) = (Dh)_{\bar{u}} \alpha'(0) = \sum_{i=1}^d c_i \frac{\partial h}{\partial u_i}(\bar{u}).$$



In the other direction, if $\alpha: \mathbb{R} \rightarrow M^d$ is any curve

$$\text{with } \alpha(0) = x, \text{ then } \alpha'(t) = [h^{-1}(\alpha)]'(t) = (Dh)_{\bar{u}} (h^{-1})'(t) \in \text{Span}\left(\frac{\partial h}{\partial u_i}(\bar{u})\right). \square$$

The picture is:



Defin: The tangent bundle to a smooth manifold $M^d \subseteq R^N$

is the union of all tangent spaces $T_x M^d$, $x \in M^d$:

$$T(M^d) = \{(x, \vec{v}) \in M^d \times R^N \mid \vec{v} \in T_x M^d\}.$$

We topologize $T(M^d)$ as a subspace of $M^d \times R^N$,

so that the projection map $\begin{matrix} \pi \\ \downarrow \end{matrix} : T(M^d) \times (x, \vec{v}) \rightarrow M^d \times X$ is continuous.

Lemma: $T(M^d)$ is locally trivial: that is, we have

homeomorphisms $\pi^{-1}(V) \xrightarrow{\cong} V \times R^d$ whenever $V \subseteq h(U)$ for some

local parametrization $h: U \rightarrow M^d$. Moreover, these homeomorphisms

are linear on each fiber, i.e. $\varphi|_{T_x(M^d)} : \{x\} \times R^d \rightarrow \{x\} \times R^d$ is linear.

[Note: φ is a homeomorphism, so linear on fibers \Rightarrow isom. on fibers.]

Proof: The parametrization h gives us a choice of basis $\left\{ \frac{\partial h}{\partial u_j}(x) \right\}_{j=1}^d$ for $T_x(M^d)$. We now define

$$\varphi\left(x, \sum c_i \frac{\partial h}{\partial u_j}(h(x))\right) = \left(x, \begin{pmatrix} c_1 \\ \vdots \\ c_d \end{pmatrix}\right) \in V \times R^d;$$

$$\text{and } \tilde{\varphi}'(x, (c_i)) = \left(x, \sum c_i \frac{\partial h}{\partial u_j}(h(x))\right).$$

These maps are well-defined and inverse to one another.

Since $\frac{\partial h}{\partial u_i}$ is continuous, we see that φ^{-1} is continuous. To check continuity of φ , we use some basic linear algebra and Kramer's Rule. We will show that the injective linear maps $(Dh)_{h^{-1}x}$ have continuously varying left inverses L_x .

We claim, then, that $\varphi(x, \vec{w}) = (x, L_x \vec{w})$, which is continuous. Indeed, we can write $\vec{w} = \sum c_i \frac{\partial h}{\partial u_i}(h^{-1}x) = (Dh)_{h^{-1}x} \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix}$ for some $\begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix}$, and then $(x, L_x \vec{w}) = (x, L_x(Dh)_{h^{-1}x} \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix}) = (x, \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix})$, which was our definition of $\varphi(x, \vec{w})$.

Now, say $A : \mathbb{R}^d \hookrightarrow \mathbb{R}^n$ is a linear injection. Let $a_i = Ae_i$ and find $\vec{a}_{d+1}, \dots, \vec{a}_n \in \mathbb{R}^n$ such that $\{\vec{a}_1, \dots, \vec{a}_n\} \subseteq \mathbb{R}^n$ is a basis

for \mathbb{R}^n . Then we have a linear map $\mathbb{R}^n \xrightarrow{L} \mathbb{R}^d$ which sends

$$\sum c_i \vec{a}_i \text{ to } \begin{pmatrix} c_1 \\ \vdots \\ c_d \end{pmatrix}, \text{ and } LA = I_d \text{ because } L(A \begin{pmatrix} c_1 \\ \vdots \\ c_d \end{pmatrix}) = L \left(\sum_{i=1}^d c_i a_i \right) = \begin{pmatrix} c_1 \\ \vdots \\ c_d \end{pmatrix}.$$

We claim that the matrix of L is

$$d \underbrace{\left\{ \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix} \cdot (\vec{a}_1 \cdots \vec{a}_n)^{-1} \right\}}_P$$

Since $(\vec{a}_1 \cdots \vec{a}_n) \vec{e}_i = \vec{a}_i$, we have $(\vec{a}_1 \cdots \vec{a}_n)^{-1} \vec{a}_i = e_i$,

and hence $P \vec{a}_i = \begin{cases} e_i, & i \leq d \\ 0, & \text{else} \end{cases}$. This agrees with our definition of $L(a_i)$, so P and L agree on a basis.

Now, the formula for P shows that P varies continuously with the vectors $\vec{a}_1, \dots, \vec{a}_n$, because Kramer's rule gives a continuous formula for the inverse of a matrix. We must show that if the vectors $\{\vec{a}_1, \dots, \vec{a}_d\}$ are varying continuously, we can choose the remaining vectors $\{\vec{a}_{d+1}, \dots, \vec{a}_n\}$ continuously.

But since $\det(\vec{a}_1 \dots \vec{a}_n) \neq 0$, there exist open neighborhoods around each $\vec{a}_1, \dots, \vec{a}_d$ so that for $\vec{a}'_1, \dots, \vec{a}'_d$ in these neighborhoods, $\det(\vec{a}'_1 \dots \vec{a}'_d \vec{a}_{d+1} \dots \vec{a}_n) \neq 0$, and hence $\{\vec{a}'_1, \dots, \vec{a}'_d, \vec{a}_{d+1}, \dots, \vec{a}_n\}$ is still a basis.

So in fact, the same choice of $\vec{a}_{d+1} \dots \vec{a}_n$ will work for nearby $\{\vec{a}_1, \dots, \vec{a}_d\}$. This allows us to choose (locally) left inverses to $(Dh)^{-1}_{\vec{x}}$ which vary continuously. This shows that φ is indeed continuous.

with well-known claims. It follows that the inverse of $[Dh]_x = [a_1'(x)] \cdots [a_N'(x)]$
 The continuity of a_i' 's has completed the proof. \square

[A man says this lemma is not fully convincing. Is there
 a better proof?]

The tangent bundle $T(M^d)$ is our first example
 of a vector bundle.

Def'n: A vector bundle $V \xrightarrow{\pi} B$ is a pair of spaces

(V, B) , together with a continuous projection map $\pi: V \rightarrow B$,

and a choice of a real vector space structure on each fiber
 $\pi^{-1}(b)$ ($b \in B$). This structure must satisfy the local
triviality condition:

For each $b \in B$ there is an open nbhd $U \subset B$
 and a homeomorphism $V|_U = \pi^{-1}(U) \xrightarrow{\cong} U \times \mathbb{R}^n$ (for some n)
 which is linear on each fiber $\pi^{-1}(b)$, $b \in U$.

Def'n A map of vector bundles $(V, B) \rightarrow (V', B')$ is a comm. diag.

$$\begin{array}{ccc} V & \xrightarrow{\varphi} & V' \\ \pi \downarrow & & \downarrow \pi' \\ B & \xrightarrow{\psi} & B' \end{array}$$

such that φ restricts to a linear map on each fiber $\pi^{-1}(b)$.


Smooth maps b/w mflds give the first examples of
 maps between vector bundles:

Def'n: A smooth map $f: M^d \rightarrow N^r$ b/w smooth mflds

is a fcn such that for each $x \in M^d$ there exist parametrizations

$$h: U_1 \rightarrow M^d, \quad g: U_2 \rightarrow N^r \quad (x \in h(U_1), \quad f(x) \in g(U_2))$$

such that $\bar{g} \circ f \circ h: U_1 \rightarrow U_2$ is smooth.



A smooth map $M^d \xrightarrow{f} N^r$ induces a map (its total derivative)

$$Df: TM^d \rightarrow TN^r$$

$$\alpha'(t) \longmapsto (f \circ \alpha)'(t)$$

$$T_{\alpha(t)} M^d \quad T_{f(\alpha(t))} N^r$$

where $\alpha: (a, b) \rightarrow M^d$ is any smooth curve. Note that

with respect to the bases $\left\{\frac{\partial h}{\partial u_j}\right\}_1^d, \left\{\frac{\partial g}{\partial v_i}\right\}_1^r$, the restriction of

Df to the fiber over $x \in M^d$ has matrix $\left(\frac{\partial(g \circ f \circ h^{-1})}{\partial u_j}(h^{-1}x)\right)_{ij}$. (IEn).

particular, Df is linear on each fiber, hence a map
of vector bundles.

A Useful Fact:

We call a map $\begin{array}{ccc} V & \xrightarrow{\varphi} & W \\ \pi \downarrow & & \downarrow \pi' \\ B & \xleftarrow{\quad \quad} & \end{array}$ of vector bundles over the same base an isomorphism if it is an isom. on each fiber.

Lemma 2.3: If φ is an isom. of v. bldls over B , then φ is a homeomorphism.

Proof: See MT+S. The main point is that Cramer's Rule shows that the map $A \mapsto A^{-1}$ is a continuous map $GL_n \mathbb{R} \rightarrow GL_n \mathbb{R}$. \square