

Lecture 12

Bundles over Paracompact Spaces

We have used the following result (to define $C_c(L_E) \subset C(E)$

for example):

Theorem: If $E \rightarrow X$ is a vector bundle over a paracompact Hausdorff space X , then there exists a diagram

$$\begin{array}{ccc} E & \rightarrow & \mathbb{R}^n \\ \downarrow & & \downarrow \\ X & \rightarrow & \text{Gr}_n(\mathbb{C}^\infty) \end{array}$$

The corresponding result holds in the real case as well.
 [Remark: MS proves this without assuming X is Hausdorff!]
Proof: First, we claim that it suffices to construct

a continuous, linear injection $E \xrightarrow{j} \mathbb{C}^\infty \cong \mathbb{C}^n \oplus \mathbb{C}^n \oplus \dots$

~~...~~

Given such a map j , we define

$$\begin{array}{ccc} E & \xrightarrow{\tilde{f}} & \mathbb{R}^n \\ \pi \downarrow & & \downarrow \\ X & \xrightarrow{f} & \text{Gr}_n(\mathbb{C}^\infty) \end{array} \quad \text{by } \tilde{f}(e) = \left(j(\text{fiber through } e), j(e) \right) \text{ and } f(x) = j(\pi^{-1}x).$$

Note that locally, f has the form

$$\begin{array}{ccc} \text{local basis} \uparrow & E|_U \cong \mathbb{R}^n \times U & \xrightarrow{j(x-x_i)} \mathbb{R}^n \oplus \mathbb{R}^n \oplus \dots \oplus \mathbb{R}^n \\ \downarrow & \downarrow & \downarrow \\ x & U & \xrightarrow{f} \text{Gr}_n(\mathbb{C}^\infty) \end{array}$$

So f is continuous.

So we must construct a map

$$E \rightarrow \mathbb{C}^\infty \cong \mathbb{C}^n \oplus \mathbb{C}^n \oplus \dots$$

Lemma: If X is paracompact, and $\{U_i\}_{i \in I}$ is an open cover of X , then there exists a countable open cover $\{V_k\}_{k=1}^\infty$ of X such that

1) Each V_k can be written as a disjoint union

$$V_k = \bigsqcup_{j \in J} V_k^j$$

with $V_k^j \subseteq U_{i(j)}$ for some $i = i(j)$

2) There is a locally finite partition of unity $\{\varphi_k\}_{k=1}^\infty$ with

$$\text{supp}(\varphi_k) \subseteq V_k$$

Assuming the Lemma, we can easily construct the desired map $j: E \rightarrow \mathbb{C}^\infty$:

Let $\{U_i\}_{i \in I}$ be a cover of X over which E is trivial, and let $\{V_k\}_{k=1}^\infty$ be the cover in the Lemma.

Note that condition 1) implies that $E|_{V_k}$ is trivial for each k . Choose trivializations $\psi_k: E|_{V_k} \rightarrow V_k \times \mathbb{C}^n$, and let π_k denote $E|_{V_k} \rightarrow V_k \times \mathbb{C}^n \xrightarrow{\pi_k} \mathbb{C}^n$. Letting φ_k denote the partition of unity in 2), we

define $j(e) = \bigoplus_{k=1}^\infty \varphi_k(\pi e) \cdot \psi_k \in \mathbb{C}^n \oplus \mathbb{C}^n \oplus \dots \cong \mathbb{C}^\infty$

Since only finitely many φ_k are non-zero at $\pi(e)$, this point lies in $\bigoplus_i \mathbb{C}^n$.

Also, j is injective b/c at each $x \in X$, some φ_k must be non-zero (b/c $\sum_{k=1}^{\infty} \varphi_k(x) = 1$ at each $x \in X$). This completes the proof of the theorem.

Proof Lemma: Since X is paracompact Hausdorff,

\exists a (locally finite) part. of \mathcal{I} subordinate to $\{U_i\}_{i \in I}$.

This means a collection of fns $\{\varphi_j\}_{j \in J}$ s.t.

- $\varphi_j: X \rightarrow \mathbb{R}_{\geq 0}$
- $\text{supp}(\varphi_j) = \varphi_j^{-1}(\mathbb{R}_{>0})$ is contained in some U_i
- For each $x \in X$, \exists an open nbhd $W \ni x$ s.t. only finitely many φ_j are non-zero on W .

Define, for each finite set $S \subseteq I$,

$$V_S = \{x \in X \mid \forall s \in S, \forall i \notin S, \varphi_s(x) > \varphi_i(x)\}.$$

Note that if $x \in X$, $\exists W \ni x$ s.t. only $\varphi_{i_1}, \dots, \varphi_{i_n}$ are

non-zero on W , so $V_S \cap W = \bigcap_{\substack{s \in S \\ j=1, \dots, n}} \{x \in W \mid \varphi_s(x) > \varphi_{i_j}(x)\}$

$$= \left[\bigcap_{\substack{s \in S \\ j=1, \dots, n}} (\varphi_s - \varphi_{i_j})^{-1}(\mathbb{R}_{>0}) \right] \cap W$$

which is a finite intersection of open sets in X . Hence

V_S is open in X for each (finite) set $S \subseteq I$.

Note that $V_S \subseteq U_i$ if $\text{supp}(\varphi_s) \subseteq U_i$ for some $s \in S$,

b/c each φ_s ($s \in S$) is positive on V_s .

4

Let $V_k = \bigcup_{|S|=k} V_S$. We claim that

$\{V_k\}_{k=1}^{\infty}$ is the desired cover of X . It's certainly

a cover, since for any $x \in X$, $x \in V_{\{s \in I \mid \varphi_s(x) > 0\}}$.

Next, we claim that $V_S \cap V_{S'} = \emptyset$ if $|S| = |S'|$. Since

$S \neq S'$, $S' \neq S$, we can choose $s \in S \setminus S'$, $s' \in S' \setminus S$.

Then any point $x \in V_S \cap V_{S'}$ would have to satisfy both

$$\begin{aligned} \varphi_s(x) &> \varphi_{s'}(x) \\ \text{and } \varphi_{s'}(x) &> \varphi_s(x), \end{aligned}$$

which is impossible.

So $V_k = \bigsqcup_{|S|=k} V_S$, and each V_S lies in some U_i .

Finally, ~~we consider~~ consider a part. of $\mathbb{1}$ sub. to $\{V_k\}_{k=1}^{\infty}$, say $\{\varphi_{\alpha}\}_{\alpha \in A}$

and let $\varphi_k = \sum \{\varphi_{\alpha} : \text{supp}(\varphi_{\alpha}) \subseteq V_k \text{ but not in } V_1, \dots, V_{k-1}\}$.

~~Then $\{\varphi_k\}_{k=1}^{\infty}$ is still locally~~

Then $\text{supp}(\varphi_k) \subseteq \bigcup \text{supp}(\varphi_{\alpha}) \subseteq V_k$; $\sum_{k=1}^{\infty} \varphi_k(x) = \sum_{\alpha \in A} \varphi_{\alpha}(x) = \mathbb{1}$;

and $\{\varphi_k\}_k$ is locally finite b/c $\{\varphi_{\alpha}\}$ was locally finite, and each φ_{α} appears as a summand in just one φ_k . \square

Some Important Facts about Char. Classes

Theorem: If X is a
then line bdlcs over X are completely determined
by their first Chern class (cplx case) or their
first Stiefel-Whitney class (real case).

Proof: We need to show that if $c_1(L) = c_1(M)$
then $L \cong M$. Let $f: X \rightarrow \mathbb{C}P^\infty$, $g: X \rightarrow \mathbb{C}P^\infty$
be classifying maps for L and M (resp.). Then
the induced maps
$$f^*, g^*: H^* \mathbb{C}P^\infty = \mathbb{Z}[\alpha] \longrightarrow H^* X$$

~~$$\alpha \longmapsto c_1(L) = c_1(M)$$~~
$$\alpha \longmapsto c_1(L) = c_1(M)$$

are completely determined by the image of α ,

so $f^* = g^*$. Hence we need to show that maps
from CW cplx into $\mathbb{C}P^\infty$ are completely determined

(up to htpy) by their effect in cohomology (w/ \mathbb{Z} -coeff's).

Theorem: If Z is a space with just one
non-zero htpy group $\pi_n(Z) = \pi$ (with π

abelian if $n=1$) then $[X, Z] \cong H^n(X; \pi)$
for any CW cplx X . unbased htpy classes of maps

[Remark: Spaces like Z , with one non-zero $h_{2p+1, 2p}$, are called Eilenberg-MacLane spaces, and are usually denoted $Z = K(\pi, n)$. Up to $h_{2p+1, 2p}$, there is a unique ~~topological space~~ CW model for $K(\pi, n)$.]

This result applies to both $CP^\infty = K(\mathbb{Z}, 2)$ and $RP^\infty = K(\mathbb{Z}/2, 1)$, b/c

• $CP^\infty = Gr(1, \mathbb{C}^\infty) = BU(1) \Rightarrow \pi_* CP^\infty = \pi_{*-1} U(1)$
 $= \begin{cases} \mathbb{Z}, & * = 2 \\ 0, & \text{else} \end{cases}$
 Note: $U(1) \cong S^1$

• $RP^\infty = Gr(1, \mathbb{R}^\infty) = BO(1) \Rightarrow \pi_* RP^\infty = \pi_{*-1} O(1)$
 $O(1) = \{\pm 1\}$
 $= \begin{cases} \mathbb{Z}/2, & * = 1 \\ 0, & \text{else} \end{cases}$

[Remark: The isomorphism $\pi_* BG \cong \pi_0 G$ is an isom. of groups.]

~~... are~~
~~The isomorphism $[X, Z] \cong \pi_* X$~~

~~Old~~ The isomorphism

$$[X, K(\pi, n)] \xrightarrow{\cong} H^n(X)$$

is given by sending $f: X \rightarrow K(\pi, n)$ to $f^*(c)$

for a particular "universal class" $c \in H^n(K(\pi, n))$. ← Cohom. w/ π -coeff's

Hence if two maps $f, g: X \rightarrow K(\pi, n)$ have the same effect in cohomology, they are homotopic. This completes the proof. \square

The universal class in $H^n(K(\pi, n); \pi)$:

This class can be described in terms of the Hurewicz map ~~old~~

$$\pi_n \mathbb{Z} \rightarrow H_n(\mathbb{Z}; \mathbb{Z})$$

$$\alpha: S^n \rightarrow \mathbb{Z} \mapsto \alpha_* [S^n]$$

↪ Fundamental class in $H_n(S^n; \mathbb{Z})$

~~Applied to $\mathbb{Z} = K(\mathbb{Z}, 1)$~~

Theorem [Hurewicz Thm]: If $\pi_k \mathbb{Z} = 0$ for $k < n$,

then the Hurewicz map $\pi_n \mathbb{Z} \rightarrow H_n(\mathbb{Z}; \mathbb{Z})$ is an isom. (When $n=1$, we must also assume $\pi_1 \mathbb{Z}$ is abelian).

8

Now if Z is a $K(\pi, n)$, we define $\ell \in H^n(K(\pi, n), \pi)$
~~to be~~ to be the image of $\text{Id}: \pi \rightarrow \pi$ under the maps
 $\text{Hom}(\pi, \pi) \cong \text{Hom}(\pi_n(K(\pi, n)), \pi) \xrightarrow{\text{Hurwicz}} \text{Hom}(H_n(K(\pi, n), \mathbb{Z}), \pi)$
 $\xrightarrow{\text{UCT}} H^n(K(\pi, n); \pi).$

This class ℓ depends only on our identification
 $\pi \cong \pi_n(K(\pi, n))$, and if we replace π by the isomorphic
 group $\pi_n(K(\pi, n))$ everywhere, ℓ becomes canonical.

For a proof that $H^n(X; \pi) \cong [X, K(\pi, n)]$
 $f^*(\ell) \leftarrow f$

is an isom., see Hatcher, Chap. 4. (Possibly
 there will be a HW exercise containing another proof.)

We have defined and studied Chern classes and
 Stiefel-Whitney classes, and we observed that
 w_i, c_i are not always zero (b/c there exist non-trivial
 line bdl's).

Theorem: The Chern classes of $\begin{matrix} \gamma_n \\ \downarrow \\ \text{Gr}_n \mathbb{C}^\infty \end{matrix}$ and the Stiefel-Wh.
 Classes of $\begin{matrix} \gamma_n \\ \downarrow \\ \text{Gr}_n \mathbb{R}^\infty \end{matrix}$ are all non-zero.

9

Pf: It suffices to show that there exist bundles
w/ W_k, c_k non-zero. We'll work in the \mathbb{C} case;
the real case is identical.

$$\text{Let } \pi_i : \underbrace{\mathbb{C}P^\infty \times \dots \times \mathbb{C}P^\infty}_k \rightarrow \mathbb{C}P^\infty$$

denote the i^{th} projection, and consider the

$$\text{bundle } \gamma_1 \times \dots \times \gamma_1 \cong \pi_1^* \gamma_1 \oplus \dots \oplus \pi_k^* \gamma_1.$$

$$\text{We have } c(\gamma_1 \times \dots \times \gamma_1) = \prod c(\gamma_1) = \prod (1 + c_i \gamma_1),$$

and we claim that each term in this sum is non-zero

i.e. each Chern class c_1, \dots, c_k of $\gamma_1 \times \dots \times \gamma_1$ is
non-zero. This follows from the Kunneth Thm,

which says that

$$H^*(\mathbb{C}P^\infty \times \dots \times \mathbb{C}P^\infty) \cong \bigotimes_{i=1}^k H^*(\mathbb{C}P^\infty)$$

$$\pi_1^* \gamma_1 \cup \dots \cup \pi_k^* \gamma_1 \longleftarrow \gamma_1 \oplus \dots \oplus \gamma_1$$

meaning that there can be no relations among the classes

$\pi_i^*(c_1 \gamma_1)$. Since the degree l term in $\prod (1 + c_i \gamma_1)$
is a poly. in $c_i \gamma_1$, it must be non-zero. \square

In fact, more is true:

Then: ~~$H^*(Gr_n(\mathbb{C}^\infty))$~~ $\cong \mathbb{Z}$ -coeffs
 $H^*(Gr_n(\mathbb{C}^\infty)) \cong \mathbb{Z}[c_1, c_2, \dots, c_n]$

where $c_i = c_i(\gamma_n)$, the Chern classes of the universal bdl, and

$$H^*(Gr_n(\mathbb{R}^\infty), \mathbb{Z}/2) \cong \mathbb{Z}/2[w_1, \dots, w_n]$$

w/ $w_i = w_i(\gamma_n)$.

This theorem says that up to multiplicative combinations, the Stiefel-Whitney / Chern classes account for all characteristic classes of vector bdl's.

Sketch of Proof (MS § 7):

We have shown that $H^*(Gr_n(\mathbb{C}^\infty; \mathbb{Z}))$ and ~~$H^*(Gr_n(\mathbb{R}^\infty; \mathbb{Z}/2)$~~ $H^*(Gr_n(\mathbb{R}^\infty; \mathbb{Z}/2))$ contain poly. algebras on $c_1, \dots, c_n / w_1, \dots, w_n$, b/c relations among these classes would imply relations among the Chern classes of γ_n , ~~γ_n~~ . MS § 5 gives a cell structure on the Grassmannians, which provides the ~~corresponding~~ corresponding upper b'd on H^* . \square