EXAMPLE 9 If |x-4| < 0.1 and |y-7| < 0.2, use the Triangle Inequality to estimate |(x + y) - 11|.

SOLUTION In order to use the given information, we use the Triangle Inequality with a = x - 4 and b = y - 7:

$$|(x + y) - 11| = |(x - 4) + (y - 7)|$$

$$\leq |x - 4| + |y - 7|$$

$$< 0.1 + 0.2 = 0.3$$

$$|(x + y) - 11| < 0.3$$

Thus

EXERCISES

1-12 Rewrite the expression without using the absolute-value symbol.

1.
$$|5-23|$$

4.
$$|\pi - 2|$$

5.
$$|\sqrt{5} - 5|$$

7.
$$|x-2|$$
 if $x < 2$

8.
$$|x-2|$$
 if $x > 2$

9.
$$|x+1|$$

10.
$$|2x-1|$$

11.
$$|x^2 + 1|$$

12.
$$|1 - 2x^2|$$

13-38 Solve the inequality in terms of intervals and illustrate the solution set on the real number line.

13.
$$2x + 7 > 3$$

14.
$$3x - 11 < 4$$

15.
$$1 - x \le 2$$

16.
$$4 - 3x \ge 6$$

17.
$$2x + 1 < 5x - 8$$

18.
$$1 + 5x > 5 - 3x$$

19.
$$-1 < 2x - 5 < 7$$

20.
$$1 < 3x + 4 \le 16$$

21.
$$0 \le 1 - x < 1$$

22.
$$-5 \le 3 - 2x \le 9$$

23.
$$4x < 2x + 1 \le 3x + 2$$

24.
$$2x - 3 < x + 4 < 3x - 2$$

25.
$$(x-1)(x-2) > 0$$

26.
$$(2x + 3)(x - 1) \ge 0$$

27.
$$2x^2 + x \le 1$$

28.
$$x^2 < 2x + 8$$

29.
$$x^2 + x + 1 > 0$$

30.
$$x^2 + x > 1$$

31.
$$x^2 < 3$$

32.
$$x^2 \ge 5$$

33.
$$x^3 - x^2 \le 0$$

34.
$$(x + 1)(x - 2)(x + 3) \ge 0$$

35.
$$x^3 > x$$

36.
$$x^3 + 3x < 4x^2$$

37.
$$\frac{1}{v} < 4$$

38.
$$-3 < \frac{1}{r} \le 1$$

39. The relationship between the Celsius and Fahrenheit temperature scales is given by $C = \frac{5}{9}(F - 32)$, where C is the temperature in degrees Celsius and F is the temperature in degrees Fahrenheit. What interval on the Celsius scale corresponds to the temperature range $50 \le F \le 95$?

- **40.** Use the relationship between C and F given in Exercise 39 to find the interval on the Fahrenheit scale corresponding to the temperature range $20 \le C \le 30$.
- 41. As dry air moves upward, it expands and in so doing cools at a rate of about 1°C for each 100-m rise, up to about 12 km.
 - (a) If the ground temperature is 20°C, write a formula for the temperature at height h.
 - (b) What range of temperature can be expected if a plane takes off and reaches a maximum height of 5 km?
- 42. If a ball is thrown upward from the top of a building 128 ft high with an initial velocity of 16 ft/s, then the height h above the ground t seconds later will be

$$h = 128 + 16t - 16t^2$$

During what time interval will the ball be at least 32 ft above the ground?

43–46 Solve the equation for x.

43.
$$|2x| = 3$$

44.
$$|3x + 5| = 1$$

45.
$$|x+3| = |2x+1|$$

46.
$$\left| \frac{2x-1}{x+1} \right| = 3$$

47-56 Solve the inequality.

47.
$$|x| < 3$$

48.
$$|x| \ge 3$$

49.
$$|x-4| < 1$$

50.
$$|x-6| < 0.1$$

51.
$$|x+5| \ge 2$$

52.
$$|x+1| \ge 3$$

53.
$$|2x - 3| \le 0.4$$

54.
$$|5x - 2| < 6$$

55.
$$1 \le |x| \le 4$$

56.
$$0 < |x - 5| < \frac{1}{2}$$

緩顯

which is in slope-intercept form with $m = -\frac{2}{3}$. Parallel lines have the same slope, so the required line has slope $-\frac{2}{3}$ and its equation in point-slope form is

$$y - 2 = -\frac{2}{3}(x - 5)$$

We can write this equation as 2x + 3y = 16.

EXAMPLE 8 Show that the lines 2x + 3y = 1 and 6x - 4y - 1 = 0 are perpendicular. SOLUTION The equations can be written as

$$y = -\frac{2}{3}x + \frac{1}{3}$$
 and $y = \frac{3}{2}x - \frac{1}{4}$

from which we see that the slopes are

$$m_1 = -\frac{2}{3}$$
 and $m_2 = \frac{3}{2}$

Since $m_1m_2 = -1$, the lines are perpendicular.

B EXERCISES

1-6 Find the distance between the points.

line

ear

2.
$$(1, -3)$$
, $(5, 7)$

6.
$$(a, b)$$
, (b, a)

7-10 Find the slope of the line through P and Q.

8.
$$P(-1, 6)$$
, $Q(4, -3)$

9.
$$P(-3, 3)$$
, $Q(-1, -6)$

10.
$$P(-1, -4)$$
, $Q(6, 0)$

- 11. Show that the triangle with vertices A(0, 2), B(-3, -1), and C(-4, 3) is isosceles.
- 12. (a) Show that the triangle with vertices A(6, -7), B(11, -3), and C(2, -2) is a right triangle using the converse of the Pythagorean Theorem.
 - (b) Use slopes to show that ABC is a right triangle.
 - (c) Find the area of the triangle.
- 13. Show that the points (-2, 9), (4, 6), (1, 0), and (-5, 3) are the vertices of a square.
- 14. (a) Show that the points A(-1, 3), B(3, 11), and C(5, 15) are collinear (lie on the same line) by showing that |AB| + |BC| = |AC|.
 - (b) Use slopes to show that A, B, and C are collinear.
- **15.** Show that A(1, 1), B(7, 4), C(5, 10), and D(-1, 7) are vertices of a parallelogram.
- **16.** Show that A(1, 1), B(11, 3), C(10, 8), and D(0, 6) are vertices of a rectangle.

17-20 Sketch the graph of the equation.

17.
$$x = 3$$

18.
$$y = -2$$

19.
$$xy = 0$$

20.
$$|y| = 1$$

21-36 Find an equation of the line that satisfies the given conditions.

- **21.** Through (2, -3), slope 6
- **22.** Through (-1, 4), slope -3
- **23.** Through (1, 7), slope $\frac{2}{3}$
- **24.** Through (-3, -5), slope $-\frac{7}{2}$
- **25.** Through (2, 1) and (1, 6)
- **26.** Through (-1, -2) and (4, 3)
- **27.** Slope 3, y-intercept -2
- **28.** Slope $\frac{2}{5}$, y-intercept 4
- **29.** *x*-intercept 1, *y*-intercept -3
- **30.** x-intercept -8, y-intercept 6
- **31.** Through (4, 5), parallel to the x-axis
- **32.** Through (4, 5), parallel to the y-axis
- **33.** Through (1, -6), parallel to the line x + 2y = 6
- **34.** y-intercept 6, parallel to the line 2x + 3y + 4 = 0
- **35.** Through (-1, -2), perpendicular to the line 2x + 5y + 8 = 0
- **36.** Through $(\frac{1}{2}, -\frac{2}{3})$, perpendicular to the line 4x 8y = 1

37-42 Find the slope and *y*-intercept of the line and draw its graph.

37.
$$x + 3y = 0$$

38.
$$2x - 5y = 0$$

39.
$$y = -2$$

40.
$$2x - 3y + 6 = 0$$

41.
$$3x - 4y = 12$$

42.
$$4x + 5y = 10$$

43-52 Sketch the region in the xy-plane.

43.
$$\{(x, y) \mid x < 0\}$$

44.
$$\{(x, y) \mid y > 0\}$$

45.
$$\{(x, y) \mid xy < 0\}$$

46.
$$\{(x, y) \mid x \ge 1 \text{ and } y < 3\}$$

47.
$$\{(x, y) \mid |x| \le 2\}$$

48.
$$\{(x, y) \mid |x| < 3 \text{ and } |y| < 2\}$$

49.
$$\{(x, y) \mid 0 \le y \le 4 \text{ and } x \le 2\}$$

50.
$$\{(x, y) \mid y > 2x - 1\}$$

51.
$$\{(x, y) \mid 1 + x \le y \le 1 - 2x\}$$

52.
$$\{(x, y) \mid -x \le y < \frac{1}{2}(x+3)\}$$

- **53.** Find a point on the y-axis that is equidistant from (5, -5) and (1, 1).
- **54.** Show that the midpoint of the line segment from $P_1(x_1, y_1)$ to $P_2(x_2, y_2)$ is

$$\left(\frac{x_1+x_2}{2},\frac{y_1+y_2}{2}\right)$$

- 55. Find the midpoint of the line segment joining the given points.(a) (1, 3) and (7, 15)(b) (-1, 6) and (8, -12)
- **56.** Find the lengths of the medians of the triangle with vertices A(1, 0), B(3, 6), and C(8, 2). (A median is a line segment from a vertex to the midpoint of the opposite side.)

- 57. Show that the lines 2x y = 4 and 6x 2y = 10 are not parallel and find their point of intersection.
- **58.** Show that the lines 3x 5y + 19 = 0 and 10x + 6y 50 = 0 are perpendicular and find their point of intersection.
- **59.** Find an equation of the perpendicular bisector of the line segment joining the points A(1, 4) and B(7, -2).
- **60.** (a) Find equations for the sides of the triangle with vertices P(1, 0), Q(3, 4), and R(-1, 6).
 - (b) Find equations for the medians of this triangle. Where do they intersect?
- 61. (a) Show that if the x- and y-intercepts of a line are nonzero numbers a and b, then the equation of the line can be put in the form

$$\frac{x}{a} + \frac{y}{b} = 1$$

This equation is called the **two-intercept form** of an equation of a line.

- (b) Use part (a) to find an equation of the line whose x-intercept is 6 and whose y-intercept is −8.
- **62.** A car leaves Detroit at 2:00 PM, traveling at a constant speed west along I-96. It passes Ann Arbor, 40 mi from Detroit, at 2:50 PM.
 - (a) Express the distance traveled in terms of the time elapsed.
 - (b) Draw the graph of the equation in part (a).
 - (c) What is the slope of this line? What does it represent?

C Graphs of Second-Degree Equations

In Appendix B we saw that a first-degree, or linear, equation Ax + By + C = 0 represents a line. In this section we discuss second-degree equations such as

$$x^{2} + y^{2} = 1$$
 $y = x^{2} + 1$ $\frac{x^{2}}{Q} + \frac{y^{2}}{A} = 1$ $x^{2} - y^{2} = 1$

which represent a circle, a parabola, an ellipse, and a hyperbola, respectively.

The graph of such an equation in x and y is the set of all points (x, y) that satisfy the equation; it gives a visual representation of the equation. Conversely, given a curve in the xy-plane, we may have to find an equation that represents it, that is, an equation satisfied by the coordinates of the points on the curve and by no other point. This is the other half of the basic principle of analytic geometry as formulated by Descartes and Fermat. The idea is that if a geometric curve can be represented by an algebraic equation, then the rules of algebra can be used to analyze the geometric problem.

Circles

As an example of this type of problem, let's find an equation of the circle with radius r and center (h, k). By definition, the circle is the set of all points P(x, y) whose distance